

Joseph Walrath, MD
Oculoplastic Surgery

MEDICATIONS TO AVOID PRIOR TO SURGERY

Please call the office with any questions ASAP to avoid cancellation of your surgery!

ASPIRIN - CONTAINING COMPOUNDS & RELATED STOP 2 WEEKS PRIOR TO SURGERY IF APPROVED BY YOUR DOCTOR

- | | | | | | |
|------------------------|----------------------|----------------------|------------------|---------------------|---------------------|
| • 4-Way Cold Tabs | • Asacol | • Carisoprodol | • Excedrin | • Micrainin | • Robaxisal |
| • 4-Way w/ Codeine | • Ascriptin | • Compound | • Fiorgen | • Midol | • Rowasa |
| • A.C.A | • Aspergum | • Choline | • Fiorinal | • Mobidin | • Roxiprin |
| • A-A Compound | • Asprimox | • Magnesium | • Fiorital | • Mobigesic | • Salflex |
| • Acetylsalicylic Acid | • Axotal | • Trisalicylate | • Gelprin | • Mono-Gesic | • Salicylic Acid |
| • Alka-Seltzer | • Azdone | • Choline Salicylate | • Goody's Powder | • Night-Time | • Salsalate |
| • Amigesic | • Azulfidine | • Damason-P | • Halfprin | • Effervescent Cold | • Sodium Salicylate |
| • Anacin | • Backache Max | • Darvon | • Isollyl | • Norwich | • Sodal compound |
| • Arthra-G | • Strength Relief | • Dipenturn | • Lanorinal | • Olsalazine | • Soma |
| • Arthriten | • Blsmatrol | • Disaclid | • Lortab ASA | • PAC | • Sulfasalazine |
| • Arthritis Foundation | • Bayer | • Doans Pills | • Magnaprin | • Pabalate | • Synalgos |
| • Products | • BC Powder | • Easprin | • products | • Panasal | • Talwin Compound |
| • Arthritis Pain | • Buffex | • Ecotrin | • Magnesium | • Pentasa | • Trigesic |
| • Formula | • Butal / ASA / Caff | • Emagin | • Salicylate | • Pepto-Bismol | • Tricosal |
| • Arthritis Strength | • Butalbital | • Empirin | • Magsal | • Percodan | • Ursinus |
| • Bufferin | • Compound | • Endodan | • Marthritic | • Pink Bismuth | • Vanquish |
| • ASA | • Cama | • Equagesic | • Mesalamine | • Relafin | |

Non-Steroidal Anti-inflammatories (NSAIDs) STOP 2 WEEKS PRIOR TO SURGERY

- | | | | | | |
|------------------|----------------|------------------|----------------|----------------------|----------------|
| • Advil | • Duract | • Lodine | • Naprelan | • Pain Reliever Tabs | • Tolmetin |
| • Aleve | • Etodolac | • Meclufenamate | • Naprosyn | • Pamprin | • Toradol |
| • Anaprox | • Feldene | • Meclomen | • Naprox X | • Piroxicam | • Vicoprofen |
| • Ansaid | • Fenoprofen | • Mefenamic Acid | • Naproxen | • Ponstel | • Voltaren |
| • Acular (drops) | • Ibuprin | • Meloxicam | • Naproxyn | • Profenal | • IBU |
| • Cataflam | • Ibuprohm | • Menadol | • Nabumetone | • Relafin | • Flurbiprofen |
| • Clinoril | • Ibuprofen | • Midol IB | • Nuprin | • Saleto | • Indochron |
| • Daypro | • Indocin | • Mobic | • Ocufen drops | • Sine-Aid IB | • Genpril |
| • Diclofenac | • Indomethacin | • Motrin | • Orudis | • Sulindac | • Haltran |
| • Difunisal | • Ketoprofen | • Nabumetone | • Oruvail | • Talwin NX | |
| • Dolobid | • Ketorolac | • Nalfon | • Oxaprozin | • Tolectin | |

Anti-Platelet / Anticoagulant Medications STOP 10 DAYS PRIOR TO SURGERY IF APPROVED BY YOUR DOCTOR

- | | | | | |
|---------------|---------------|-----------|-------------|---------------|
| • Anisindione | • Clopidogrel | • Miradon | • Plavix | • Ticlid |
| • Cilostazol | • Effient | • Pletal | • Plasugrel | • Ticlopidine |

Anticoagulant Medications STOP 5 DAYS PRIOR TO SURGERY IF APPROVED BY YOUR DOCTOR

- | | | | |
|--------------|-----------------|------------|------------|
| • Dabigatran | • Pradaxa | • Sofarin | • Warfarin |
| • Arixtra | • Fondaparineux | • Coumadin | |

Anticoagulant Medications STOP 24 HOURS PRIOR TO SURGERY IF APPROVED BY YOUR DOCTOR

- | | | | |
|--------------|--------------|-------------|------------|
| • Argatroban | • Enoxaparin | • Heparin | • Lovenox |
| • Dalteparin | • Fragmin | • Lepirudin | • Refludan |

Other Medicines to stop 2 weeks prior to surgery

- | | | | | |
|------------------|------------------|----------------------|-----------------------|------------------|
| • Anisindione | • Duragesic | • Kaodene | • Prednisone | • Scot-Tussin |
| • Cilostazol | • Contac | • Macrochantin | • Virbamycin | • Sine-off |
| • Acutrim | • Ephedra | • Methocarbamol | • Propoxyphene | • Sinex |
| • Actifed | • Cortisone | • Oxycodone | • Phenylpropanolamine | • Sinutab |
| • Actron | • Medications | • Norgestic products | • mine | • Sulfinpyrazone |
| • Anturane | • Hydrocortisone | • Orphengesic | • Pyroxate | • Triaminicin |
| • Dimetapp Sinus | • Flagyl | • products | • Ru-Tuss | • Tussanil DH |
| • Dristan | • Doxycycline | • Phenaphen/ | • Tussirex | |
| • Cheracol | • Furadantin | • Codeine #3 | • Tenuate | |

**HERBAL MEDICINES, NONTRADITIONAL MEDICINES,
VITAMINS, HOMEOPATHIC MEDICINES, DIETARY
SUPPLEMENTS: STOP ALL FOR 2 WEEKS PRIOR TO SURGERY**